

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/52, 5/10	A1	(11) International Publication Number: WO 97/08324 (43) International Publication Date: 6 March 1997 (06.03.97)
(21) International Application Number: PCT/US96/13627 (22) International Filing Date: 22 August 1996 (22.08.96) (30) Priority Data: 60/002,621 22 August 1995 (22.08.95) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventor: LAU, Allan, S.; University of California, Dept. of Pediatrics, Room 6E6, SFGH, 1001 Potrero Avenue, San Francisco, CA 94110 (US). (74) Agent: BERLINER, Robert; Robbins, Berliner & Carson, 5th floor, 201 North Figueroa Street, Los Angeles, CA 90012-2628 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHODS FOR ENHANCING THE PRODUCTION OF INTERFERON IN CELL CULTURE (57) Abstract Methods for enhancing the production of interferons in animal cell culture are described. These methods rely on the manipulation of the cellular levels of certain inducers of interferon production, in particular cellular levels of double-stranded-RNA-dependent kinase (dsRNA-PKR, or PKR). In cell cultures that overproduce PKR, interferon synthesis is induced to high levels, and significant amount of interferon can be recovered without conventional induction of interferon by virus.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

1.

METHODS FOR ENHANCING THE PRODUCTION OF INTERFERON IN CELL CULTURE

5

INTRODUCTION

Technical Field

The present invention relates to methods for enhancing the production of interferon in cell culture.

10

Background

Interferons (IFNs) can be classified into two major groups based on their primary sequence. Type I interferons, IFN- α and IFN- β , are encoded by a super family of intron-less genes consisting of the IFN- α gene family and a single IFN- β gene. Type II interferon, or IFN- γ , consists of only a single type and is restricted to lymphocytes (T-cells and natural killer cells). Type I interferons mediate diverse biological processes including induction of antiviral activities, regulation of cellular growth and differentiation, and modulation of immune functions (Sen, G. C. & Lengyel, P. (1992) *J. Biol. Chem.* 267, 5017-5020; Pestka, S. & Langer, J. A. (1987) *Ann. Rev. Biochem.* 56, 727-777). The induced expression of Type I IFNs, which include the IFN- α and IFN- β gene families, is detected typically following viral infections. Many studies have identified promoter elements and transcription factors involved in regulating the expression of Type I IFNs (Du, W., Thanos, D. & Maniatis, T. (1993) *Cell* 74, 887-898; Matsuyama, T., Kimura, T., Kitagawa, M., Pfeffer, K., Kawakami, T., Watanabe, N., Kundig, T. M., Amakawa, R., Kishihara, K., Wakeham, A., Potter, J., Furlonger, C. L., Narendran, A., Suzuki, H., Ohashi, P. S., Paige, C. J., Taniguchi, T. & Mak, T. W. (1993) *Cell* 75, 83-97; Tanaka, N. & Taniguchi, T. (1992) *Adv. Immunol.* 52, 263-81). However, it remains unclear what are the biochemical cues that signify viral infections to the cell and the signaling mechanisms involved (for a recent

30

2.

review of the interferon system, see Jaramillo et al. *Cancer Investigation* 1995 13:327-337).

IFNs belong to a class of negative growth factors having the ability to inhibit growth of a wide variety of cells with both normal and transformed phenotypes. IFN therapy has been shown to be beneficial in the treatment of human malignancies such as Kaposi's sarcoma, chronic myelogenous leukemia, non-Hodgkin's lymphoma and hairy cell leukemia as well as the treatment of infectious diseases such as papilloma virus (genital warts) and hepatitis B and C (reviewed by Gutterman *Proc. Natl Acad Sci.* 91:1198-1205 1994). Recently genetically engineered, bacterially produced IFN- β was approved for treatment of multiple sclerosis, a relatively common neurological disease affecting at least 250,000 patients in the United States alone.

Currently, IFNs for therapeutic use are produced from two major sources, natural IFNs from human leukocytes or leukocyte cell lines and recombinant IFNs produced in bacterial cells. Natural IFNs are considered to be superior as they consist of the entire complement of IFNs and have the proper structure, but they are expensive and time-consuming to produce. Bacterially produced recombinant IFN is cheaper and more efficient to make, but studies have shown much higher rates of rejection for the bacterially produced protein, particularly after long term usage. For instance, previous medical studies have shown that the incidence of rejection as reflected by antibody formation can be as high as 20 to 38% for bacterially-produced IFN compared with only 1.2% for natural IFN- α (Antonelli et al. *J. Inf. Disease* 163:882-885 1991; Quesada et al. *J. Clin. Oncology* 3:1522-1528 1985). Thus, a method for enhancing the production of natural IFN to make it less expensive to produce would be advantageous.

IFNs elicit their biological activities by binding to their cognate receptors followed by signal transduction leading to induction of IFN-stimulated genes (ISGs). Some ISGs have been characterized and their biological activities examined. The best studied examples of ISGs include a double-stranded-RNA-dependent kinase (dsRNA-PKR, or just PKR, formerly known as p68 kinase), 2'-5'-linked oligoadenylate (2-5A) synthetase, and Mx proteins (Taylor JL,

3.

Grossberg SE. *Virus Research* 1990 15:1-26.; Williams BRG. *Eur. J. Biochem.* 1991 200:1-11.).

PKR (short for protein kinase, RNA-dependent) is the only identified dsRNA-binding protein known to possess a kinase activity. PKR is a serine/threonine kinase whose enzymatic activation requires dsRNA binding and consequent autophosphorylation (Galabru, J. & Hovanessian, A. (1987) *J. Biol. Chem.* 262, 15538-15544; Meurs, E., Chong, K., Galabru, J., Thomas, N. S., Kerr, I. M., Williams, B. R. G. & Hovanessian, A. G. (1990) *Cell* 62, 379-390). PKR has also been referred to in the literature as dsRNA-activated protein kinase, P1/e1F2 kinase, DAI or dsl (for dsRNA-activated inhibitor), and p68 (human) or p65 (murine) kinase. Analogous enzymes have been described in rabbit reticulocytes, different murine tissues, and human peripheral blood mononuclear cells (Farrel et al. (1977) *Cell* 11, 187-200; Levin et al. (1978) *Proc. Natl Acad. Sci. USA* 75, 1121-1125; Hovanessian (1980) *Biochimie* 62, 775-778; Krust et al. (1982) *Virology* 120, 240-246; Buffet-Janvresse et al. (1986) *J. Interferon Res.* 6, 85-96). The best characterized *in vivo* substrate of PKR is the alpha subunit of eukaryotic initiation factor-2 (eIF-2a) which, once phosphorylated, leads ultimately to inhibition of cellular and viral protein synthesis (Hershey, J. W. B. (1991) *Ann. Rev. Biochem.* 60, 717-755). This particular function of PKR has been suggested as one of the mechanisms responsible for mediating the antiviral and antiproliferative activities of IFN- α and IFN- β . An additional biological function for PKR is its putative role as a signal transducer. Kumar *et al.* demonstrated that PKR can phosphorylate I κ B α , resulting in the release and activation of nuclear factor κ B (NF- κ B) (Kumar, A., Haque, J., Lacoste, J., Hiscott, J. & Williams, B. R. G. (1994) *Proc. Natl. Acad. Sci. USA* 91, 6288-6292). Given the well-characterized NF- κ B site in the IFN- β promoter, this may represent a mechanism through which PKR mediates dsRNA activation of IFN- β transcription (Visvanathan, K. V. & Goodbourne, S. (1989) *EMBO J.* 8, 1129-1138).

The present inventor have surprisingly discovered that manipulating the expression of certain ISGs can have beneficial uses in interferon production. They have discovered that over-expression of the PKR protein induces overproduction

4.

of the IFN- α and IFN- β interferons, which is useful for the enhanced production of interferon in animal cell culture.

Relevant Literature

- 5 Currently there are two major approaches to large-scale production of interferons: recombinant IFN produced in bacterial or mammalian cells or natural IFNs from human leukocyte cells following stimulation with viruses or other IFN inducers. U.S. Patent No. 5,376,567 and U.S. Patent No. 4,966,843 describe the production in Chinese hamster ovary cells of a recombinant human interferon; U.S.
- 10 Patent No. 5,196,323 describes the production of recombinant human IFN- α in E. coli cells. A number of patents describe the production of interferon from human leukocyte cells using a variety of protocols; for example, U.S. Patent No. 4,745,053 describes a process for producing interferon from whole human blood using a viral inducer, U.S. Patent No. 4,680,261 describes a process for inducing
- 15 production of interferon in mammalian cell culture using an ascorbic acid derivative or an inorganic vanadium compound, and U.S. Patent No. 4,548,900 describes a process for the induction of interferon using a polyhydric alcohol in a priming stage. The major disadvantage of the current methods of interferon production is that typically virus is used as the IFN inducer because other inducers
- 20 do not produce high enough levels of interferon for most commercial purposes. The virus must then be removed from the interferon before use, which adds time and cost to the production method. In addition, use of virus as an inducer ultimately results in the death of the interferon-producing cells, so that no recycling and re-use of the cells is possible.
- 25 The present invention overcomes these problems by providing a interferon-production system that does not require the use of a viral inducer in order to achieve high levels of interferon production. Although viral inducers can be used with the systems of the invention, other inducers that do require removal prior to use of the IFNs are still capable of producing IFNs at commercially acceptable
- 30 levels.

5.

SUMMARY OF THE INVENTION

It is an object of this invention to provide a method for enhancing the production of interferon in animal cell culture without requiring the addition of virus as an inducer.

5 This object is generally accomplished by providing animal cell cultures in which the expression of the interferon genes is substantially increased from the normal level of expression. This may be effected by manipulating the level of expression of factors that function *in vivo* to regulate the interferon level, including interferon transcriptional regulators (for example, IRF1), interferon receptors, and
10 interferon stimulated gene products (for example PKR and 2-5A synthetase).

Thus, increased production of INF and other objects of the invention as will hereinafter become apparent are accomplished by carrying out interferon production in animal cell cultures in which the level of interferon-regulating protein activity, particularly for double-stranded-RNA-dependent kinase (PKR) and 2'-5'
15 oligoadenylate synthetase (2-5A synthetase), is significantly increased from normal levels.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Diagrammatic drawings of pCMV-PKR and pMT-PKR.

20 1(A) Diagram of the pCMV-PKR showing the human PKR cDNA in the sense orientation driven by the promoter sequence from the immediate early gene of human cytomegalovirus.

1(B) Diagram of the pMT-PKR showing the human PKR cDNA in the sense orientation driven by the cadmium inducible metallothionein II promoter.

25

Figure 2. Interferon induction in U937-PKR+ cells and control U937-neo cells.

U937-PKR+ cells and U937-neo cells ($0.5 \times 10^6/\text{ml}$) were cultured in RPMI 1640 medium, supplemented with 5% fetal bovine serum. The cells were pretreated ("primed") with or without 10 nM PMA for 20 hrs followed by induction with 10
30 $\mu\text{g}/\text{ml}$ poly r(I):poly r(C) or with 0.1 TCID₅₀/cell EMCV for 20 hrs. Twenty-four hours after induction, culture supernatants were collected and assayed for IFN- α .

6.

The open bars indicate U937-neo cells, the hatched bars indicate U937-PKR+ cells. The experimental groups indicate different treatments: 1- no treatment; 2- poly r(I):poly r(C) induction alone; 3- EMCV induction alone; 4- PMA priming alone; 5- PMA priming and poly r(I):poly r(C) induction; 6- PMA priming and EMCV induction.

DESCRIPTION OF SPECIFIC EMBODIMENTS

The present invention arose in part from the discovery by the inventor that the level of interferon production in cell culture can be regulated by control of the expression or activity of certain proteins that normally regulate interferon expression *in vivo*. These factors include interferon-specific transcriptional regulators (particularly IRF1), interferon receptors, and the gene products of certain interferon simulated genes (also called interferon-mediated antiviral responses), particularly PKR. Enhancing the expression or activity of any of these factors will result in a higher than normal level of expression of interferon genes. One result of this higher than normal expression of interferon genes is that interferon production can be achieved with greater efficiency and lower cost. The remainder of this specification will be exemplified by reference to PKR, but it is to be understood that other interferon enhancing factors as described herein can be used in place of PKR.

By increasing the level of PKR protein (and therefore PKR activity) in an animal cell, interferon production can be increased. Animal cell cultures which express a higher constitutive level of PKR or in which PKR expression can be induced to higher levels are therefore useful for the production of interferons. Problems typically associated with production of interferons in cell culture, for example low yield and contamination with the virus used to induce interferon expression, are eliminated in the method of the present invention. Thus, a high yield of interferon protein can be achieved without the need for induction with virus (although virus can be used for even higher INF production).

The method relies on the use of PKR-overproducing cells as the source of the interferon, but any known or later-discovered method of interferon production

7.

may be used. No particular method of production of interferon is required except that typically a non-viral interferon inducer is used. In particular, the method comprises (a) culturing an animal cell culture capable of overproduction of PKR or an analog or homolog of PKR under conditions sufficient to overproduce PKR, and (b) treating the cell culture as appropriate to induce the expression of the interferon genes, particularly the IFN- β and IFN- α . These steps are generally followed by purifying the interferon produced. In some cases, the overproduction is not of PKR itself, but of an analog of PKR, by which is meant a non-natural protein kinase that can mediate dsRNA activation of interferon transcription (usually obtained by genetic manipulation of the gene encoding a particular PKR from a selected animal cell line). The cell culture used to produce interferon can overproduce PKR from any animal cell line used to produce the interferon, such as the PKR normally found in rabbit reticulocytes, various mouse tissues, or human peripheral blood mononuclear cells. Usually, the natural PKR for a particular cell line (as opposed to an exogenous PKR) is used for overexpression. Preferably murine p65 kinase and most preferably human p68 kinase is overproduced, in a corresponding murine or human cell culture, respectively.

Animal cell cultures capable of overproduction of a PKR gene may be isolated by any number of methods, including many that are well known in the art. Such methods include selection for cells expressing higher PKR levels, transfection with a vector containing a PKR gene (or cDNA) under control of a constitutive promoter (for example, a CMV, RSV or SV40 promoter), or transfection with a vector containing a PKR gene or cDNA under the control of an inducible promoter (for example, a heat shock promoter, a metallothionein promoter or a glucocorticoid promoter). Transfection is carried out as described previously and transfectants are selected for overproduction of PKR. By overproduction of PKR is meant higher than normal levels of PKR activity. Normal refers to under normal conditions for the cell line being used (see, for example, suggested culture conditions as supplied in catalogues for commercial cells, especially the ATCC Catalogue of Strains, available from the American Type Culture Collection, Rockville, Maryland, USA, or in scientific publications describing the cell line

8.

being used, if it is not commercially available), and in various preferred embodiments refers to production of INF under otherwise identical conditions in the parental cell line from which a production cell line is selected or produced by genetic engineering. Higher than normal level preferably means at least 150%,
5 more preferably at least 200 or 300%, most preferably at least 500%, of normal PKR level. The PKR-overproducing cell culture may be constitutive for PKR overproduction or inducible for PKR overproduction, depending on the particular method used to isolate or prepare the culture. Preferably the PKR-overproducing cell culture will be inducible for PKR overproduction in order to regulate the level
10 of PKR available for IFN induction. It will be apparent that if the PKR-overproducing cell culture is inducible for overproduction, PKR activity for overproduction will be assayed under inducing conditions. PKR activity can be determined by any of the methods known in the art or described in the following examples.

15 Any of a number of known cell cultures are useful as a parental strain for making a PKR-overproducing cell culture. Any cells normally capable of producing interferon are suitable as the parental strain, particularly cells derived from fibroblasts or immune cells, including B cells and monocytes. Particularly suitable cell cultures are pro-monocytic U937 cells, Namalwa (lymphoblastoid B)
20 cells and MRC-5 (human fibroblast) cells. Also suitable are WI-38 cells, Flow 1000 cells, Flow 4000 cells, FS-4 and FS-7 cells, MG-63 cells, CCRF-SB cells and CCRF-CEM cells.

Production of interferon in the PKR-overproducing cell culture is accomplished by methods that are well known in the art. Generally, the interferon
25 producing cells are cultured in any suitable medium, treated with an inducer to induce expression of the interferons (and with an inducer of the PKR gene, in PKR is provided under the control of an inducible promotor), and incubated following induction of interferon production. The interferon is then isolated. The cells may be primed prior to induction by addition of a priming agent. Interferon inducers
30 are many and well-known in the art and almost any known interferon inducer can be used in the present invention. Typical inducers include poly(I):poly(C), Sendai

9.

virus, Newcastle disease virus, concanavaline A, chlamydia, rickettsia, mitogen, and lipopolysaccharides. Preferably, the inducer will be a non-viral inducer. More preferably, the inducer will be poly(I):poly(C) or poly r(I):poly r(C). Non-viral inducers are preferred because the cells do not suffer the deleterious consequences that exposure to the virus provides and may therefore be recycled for re-use resulting in a lower cost of production. Typical priming agents include phorbol myristate acetate, calcium ionophores and interferons.

IFNs are purified by standard techniques that are well known in the art. These include antibody-affinity column chromatography, ion exchange chromatography, protein precipitation and centrifugation (U.S. patent No. 5,391,713); chromatography using CM-agarose, con A-agarose, and phenyl-agarose (U.S. patent No. 4,658,017); three step chromatography with glass sorbent column (U.S. patent No. 4,732,683); immunoaffinity chromatography, reverse phase HPLC, cation exchange column, and gel filtration (U.S. patent No. 4,765,903); guanidine HCl as solvent and HIC column chromatography (U.S. patent No. 4,845,032).

The levels of IFN- α in the preparations can be determined by titrating for antiviral activity on L929 (murine fibroblast) cells against National Institute of Allergy and Infectious Diseases reference standard G-023-901-527 (Lau et al. J. Clin. Invest. 82:1415-1421 1988). In this technique, 5×10^4 L929 cells are seeded into each well of 96-well microtiter plates and then incubated with two-fold serial dilutions of culture supernatants for 16 hr. The cells are subsequently challenged with EMCV or Sendai virus (0.1 or 0.01 TCID per cell). Virus-induced cytopathic effects were assessed by microscopic examination and by staining cells with 0.1% crystal violet in 5% ethanol solutions. The IFN titer is defined as the reciprocal of the highest dilution of culture supernatants capable of protecting 50% of the cells from viral-induced cytopathic effects.

Specific examples of the steps described above are set forth in the following examples. However, it will be apparent to one of ordinary skill in the art that many modifications are possible and that the examples are provided for purposes of illustration only and are not limiting of the invention unless so specified.

10.

EXAMPLES

Example 1: Preparation of constitutive PKR overexpressing cell culture

Plasmid pCMV-PKR was prepared from pRC/CMV by inserting the PKR cDNA from pBS-8.6R into the HindIII site of the pRC/CMV vector so that
5 expression of the PKR coding sequence is under control of the CMV promoter. The pRC/CMV plasmid (Invitrogen) is commonly used for eukaryotic expression. The vector offers the following features suitable for PKR transcription: i) promoter sequences from the immediate early gene of the human CMV (cytomegalovirus) for high level transcription; ii) polyadenylation signal and transcription termination
10 sequences from bovine growth hormone (BGH) gene to enhance RNA stability; iii) SV40 origin for episomal replication and simple vector rescue; iv) T7 and Sp6 RNA promoters flanking the multiple cloning site for in vitro transcription of sense and antisense RNA; and v) the ampicillin resistance gene (AMP) and ColE1 origin for selection and maintenance in *E. coli*. The vector also contains a G418
15 resistance marker (NEO) to allow for selection and identification of the plasmids after transfer to eukaryotic cells. The structure of pCMV-PKR is shown in Figure 1A.

Stable transfectants were obtained by electroporation of 5×10^6 exponentially growing U937 cells with 10 mg of each plasmid, in serum-free
20 RPMI-1640 containing DEAE-dextran (50 mg/mL), with a Gene Pulser apparatus (BioRad) set at 500 μ F, 250V. Bulk populations of stable transfectants were obtained by selection with 400 μ g/mL geneticin (GIBCO-BRL) for 3 weeks. Clonal lines were subsequently obtained by limiting dilution cloning. Cell lines were cultured in RPMI-1640 containing 10% fetal calf serum (complete media) and
25 geneticin. One representative transfectant cell line was selected and designated "U937-PKR+". The level of PKR produced in U937-PKR+ was analyzed and found to be increased approximately five-fold over normal (parental) levels. As a control, U937 cells were transfected with pRC/CMV and a representative transfectant cell line was isolated and designated "U937-neo". The transfectant cell
30 lines were tested for IFN production in the presence or absence of priming agents and/or inducers.

11.

U937-PKR+ cells and U937-neo cells were cultured in RPMI 1640 medium, supplemented with fetal bovine serum. The cells were pretreated with or without phorbol myristate acetate (PMA) (10nM) for 20 hr and with or without subsequent stimulation by poly r(I):r(C), 10 ug/ml, for 20 hr. Twenty-four hr
5 after the stimulation with poly r(I):r(C), culture supernatants were collected for IFN assays.

As shown in Figure 2, neither transfectant cell line produced IFN spontaneously in the absence of priming agents or inducers. In response to poly r(I):poly r(C), control U937-neo cells showed minimal synthesis of IFN- α either
10 with or without PMA priming (open bar 2 and 5). In contrast, U937-PKR+ cells showed increased IFN production in response to poly r(I):poly r(C) induction, with or without PMA priming, reaching a level of 4000 U/ml when treated with both poly r(I):poly r(C) and PMA (hatched bars 2 and 5). In the absence of PMA priming, EMCV induced high levels of IFN in both transfectant cell lines (more
15 than 1000 U/ml), with slightly higher levels in the U937-PKR+ cells (Bar 3). In the absence of EMCV or poly r(I):poly r(C) inducer, U937-PKR+ cells produced higher levels of IFNs than did U937-neo cells in response to PMA priming alone (Bar 4). With PMA priming and EMCV induction, both transfectant cell lines
20 produced more than 4000 U/ml IFN (Bar 6). Taken together, these results indicated that following priming with PMA, the U937-PKR+ cells are as efficient in IFN production when poly r(I):poly r(C) is used as inducer as are control cells when virus is used as inducer. Therefore the use of live virus for IFN production can be eliminated when PKR-overproducing cells are used.

Following induction by poly r(I):poly r(C), the U937-PKR+ cells were
25 examined for viability using a trypan blue exclusion assay. We found that more than 95% of the cells were viable. In contrast, the use of EMCV as an inducer resulted in the death of all of the cells. Thus, the U937-PKR+ cells can be recycled for continuous production of IFN.

12.

Example 2: Preparation of inducible PKR overexpressing cell culture

Plasmid pMT-PKR was made by replacing the CMV promoter upstream from the PKR cloning site in pCMV-PKR with a cadmium-inducible metallothionein II promoter (Hewison et al. J. Immunol. 153:5709-5719 1994).

- 5 The structure of pMT-PKR is shown in Figure 1B. Stable transfectants of pMT-PKR into U937 cells were isolated as described in Example 1 for pCMV-PKR transfectants. The MTII promoter is inducible by zinc or cadmium ions. PKR activity levels in the transfectants were measured after induction of the transfected PKR gene by addition of 20 μ M cadmium chloride.

10

All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

- 15 The invention now being fully described, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the described invention.

13.

WHAT IS CLAIMED IS:

1. A method for producing interferon in an animal cell culture, comprising:
 - (a) culturing an animal cell culture under conditions sufficient to
5 overproduce PKR; and
 - (b) treating said culture to induce interferon production.
2. A method for producing interferon in an animal cell culture, comprising:
 - (a) cultivating an animal cell culture under conditions sufficient to
10 overproduce an analog of PKR; and
 - (c) treating said culture to induce interferon production.
3. The method of Claim 2, wherein said analog is obtainable from mouse tissue.
- 15 4. The method of Claim 3, wherein said analog is murine p65 kinase.
5. The method of Claim 2, wherein said analog is obtainable from rabbit reticulocytes.
- 20 6. The method of Claim 2, wherein said analog is obtainable from human peripheral blood mononuclear cells.
7. The method of Claim 1, wherein said PKR is human p68 kinase.
- 25 8. The method of Claim 1, wherein said interferon production is induced with a non-viral inducer.
9. The method of Claim 8, wherein said non-viral inducer is poly (I):Poly(C) or poly r(I):poly r(C).

14.

10. The method of Claim 1, wherein said step (a) further comprises treating with a priming agent.

11. The method of Claim 1, further comprising the step of purifying the interferon
5 produced.

12. The method of Claim 1, wherein said animal cell culture is derived from parental cells normally capable of producing interferon.

10 13. The method of Claim 12, wherein said overproduction of PKR is achieved by transfection of said parental cells with a vector containing DNA encoding PKR under control of a promoter.

14. The method of claim 13, wherein said promoter is an inducible promoter.
15

15. The method of claim 14, wherein said inducible promoter is a metallothionein promoter.

16. The method of Claim 12, wherein said parental cells are U937 cells.

1/2

FIG. 1A





FIG. 1B

2/2




FIG. 2

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13627

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/52, 5/10

US CL : 435/69.51, 70.5

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.51, 70.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet:

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PROUD, C.G. PKR: a new name and new roles. Trends Biochem. Sci., June 1995, Vol. 20., No. 6, pages 241-246.	1-16
A	KUMAR, A., et al. Double-stranded RNA-dependent protein kinase activates transcription factor NF- κ B by phosphorylating I κ B. Proc. Natl. Acad. Sci. USA, July 1994, Vol. 91, pages 6288-6292.	1-16
A	BARBER, G.N., et al. The 58-kilodalton inhibitor of the interferon-induced double-stranded RNA-activated protein kinase is a tetratricopeptide repeat protein with oncogenic properties. Proc. Natl. Acad. Sci. USA, May 1994, Vol. 91, pages 4278-4282.	1-16

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"Z" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

13 NOVEMBER 1996

Date of mailing of the international search report

26 NOV 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DAVID L. FITZGERALD

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13627

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	DER, S.D., et al. Involvement of the double-stranded RNA-dependent kinase PKR in interferon expression and interferon-mediated antiviral activity. Proc. Natl. Acad. Sci. USA, September 1995, Vol. 92, pages 8841-8845.	1-16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/13627

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

Medline: search terms = inventors' names

SciSearch: search terms = articles cited in DER et al., Proc. Natl. Acad. Sci. USA, September 1995, cited infra.

USPTO-APS, EPO online, Derwent WPI: search terms = PKR, RNA, kinase, interferon, IFN

HPS Trailer Page
for

EAST

UserID: DRomeo_Job_1_of_1

Printer: cm1_10e14_gbekptr

Summary

<u>Document</u>	<u>Pages</u>	<u>Printed</u>	<u>Missed</u>	<u>Copies</u>
US005362490	19	19	0	1
US004474754	54	54	0	1
WO009708324	21	21	0	1
Total (3)	94	94	0	-